

**“Management Of Medically Compromised Patients In Pediatric  
Dentistry: A Literature Review”**

**Abstract:**

Lifestyle changes, dietary shifts, and other factors have contributed to an uptick in disease and disorder incidence alongside the development of new technologies and medical treatments. Because many dental operations can be stressful for patients, it's important to understand their health so that any potential adverse effects can be mitigated. This is especially true for patients whose health is already precarious. Dental implants, tooth extractions, and jaw surgeries all carry the danger of bleeding, which can be fatal for patients with bleeding problems; an overdose of laxatives can bring on an epileptic seizure; and many other complications that necessitate careful medical and dental management. It is the responsibility of the dentist to be familiar with these disorders and to know how to manage them in the dental operator; in this article, we will provide a quick overview of dental management in medically impaired patients.

Key words: Oral Cavity, Oral Manifestations, Systemic Diseases

**المخلص:**

ساهمت التغييرات في نمط الحياة والتغييرات الغذائية وعوامل أخرى في زيادة انتشار المرض والاضطراب إلى جانب تطوير تقنيات جديدة وعلاجات طبية. نظراً لأن العديد من عمليات طب الأسنان يمكن أن تكون مرهقة للمرضى، فمن المهم أن نفهم صحتهم بحيث يمكن التخفيف من أي آثار ضارة محتملة. هذا ينطبق بشكل خاص على المرضى الذين تكون صحتهم بالفعل محفوفة بالمخاطر. تتطوي عمليات زرع الأسنان وقلع الأسنان وجراحات الفك على خطر حدوث نزيف يمكن أن يكون قاتلاً للمرضى الذين يعانون من مشاكل النزيف. جرعة زائدة من المسهلات يمكن أن تسبب نوبة صرع. والعديد من المضاعفات الأخرى التي تتطلب إدارة طبية وعلاج الأسنان بعناية. تقع على عاتق طبيب الأسنان مسؤولية التعرف على هذه الاضطرابات ومعرفة كيفية إدارتها في جراحة الأسنان؛ في هذه المقالة، سوف نقدم نظرة عامة سريعة على إدارة الأسنان للمرضى المعرضين للخطر طبياً في طب أسنان الأطفال.

الكلمات المفتاحية: تجويف الفم ، مظاهر الفم ، أمراض جهازية.

### Introduction:

In both dentistry and medicine, considerable strides have been made in the previous few years, leading to more accurate diagnosis and more effective treatment of disease. Yet, the disease rate has also increased in today's population, maybe as a result of environmental changes, the use of far more chemicals, etc., as well as changes in people's lifestyles, such as food and inactivity. Not only have dental procedures and technology advanced significantly over the past decade or so, but the types of patients seen on a regular basis have also shifted significantly. Several chronic illnesses or their treatments call for adjustments to dental management, and failing to make the necessary adjustments might have substantial clinical repercussions. Assessing the patient's current health status and past medical history is crucial for dental therapy of a medically challenged patient.

A person's mouth can be used as a sentinel or early warning system, reflecting their overall state of health or illness. The mouth, as the body's primary entry point, is always under attack from pathogens such as bacteria, viruses, parasites, and fungus. Oral lesions and ulcers can appear anywhere in the mouth, including the oral mucosa, tongue, gums, extraoral skin, and other associated structures, as a result of a wide variety of systemic disorders. These oral manifestations necessitate a correct diagnosis and treatment strategy. <sup>1</sup> The oral symptoms of a child's underlying disease raise the child's risk of developing an oral disease. The oral cavity may also experience negative effects and a reduction in host resistance as a result of the treatment or drug. It is important to address dental disease in children with medical conditions since it can have serious consequences for their overall health and quality of life if left untreated. A severe dental infection might be fatal under some circumstances.

Infections in the teeth might compromise the effectiveness of medical treatment. Their dental care is crucial because of the potential danger to their health or even their lives. Children with medical conditions should be given priority for early and regular preventative dental care. Maintaining good oral hygiene practises is crucial in avoiding oral infections. When it comes to children with preexisting medical conditions, however, the need of oral disease prevention increases dramatically. Maintaining good oral hygiene is critical to the overall health of children with medical conditions, and it plays a role in the success of many procedures performed to rectify congenital abnormalities and transplant organs. Nevertheless, neglect and poor dental hygiene might have a negative impact on recovery time and success rates after surgical treatments.

### Definition of medically compromised patient:

A patient with a medical condition who is also receiving treatment for another pathology may experience further deterioration of their health.



### Discussion:

The analysis zeroes in on a select number of medical issues that dentists may face in their daily practise that need for further training and attention to avoid complications that could lead to avoidable mortality and morbidity. Diabetes mellitus, heart abnormalities, blood disorders like hemophilia, kidney disorders, neurological disorders like epilepsy, respiratory issues, infectious diseases like HIV or hepatitis, and many other diseases and disorders, as well as many drug interactions, are all included.

Obtaining a complete medical history and understanding the seriousness of the disease that may be endorsed by the patient are crucial in managing the patient with medical difficulties; additionally, each medical condition detected might impact dental care in a different way. Complication management strategies are described for a selection of medical disorders below:

- Diabetes mellitus:

Type 2 diabetes (DM) is brought on by either a lack of insulin production or an inability of insulin receptors in end organs to respond to insulin. Type 1 (insulin-dependent) and type 2 (non-insulin-dependent) diabetes are the two most common classifications of diabetes. In most cases, the onset of type 1 diabetes occurs in young adulthood or during infancy. In managing a diabetic patient, preventing hypoglycemia episodes and keeping blood glucose levels as close to normal as possible are the primary goals. Steroids and aspirin should be avoided, and oral infections should be managed. Non-surgical procedures are most effective when performed right after breakfast, followed by the regular dosing of diabetes medication. Blood glucose levels should be kept between 120 and 180 mg/dl prior to surgical procedures. Potentially detrimental effects of hyperglycemia include impaired wound healing and other dysfunctionality. Glucose levels during intravenous glucose infusion should be monitored so that soluble insulin can be administered as needed to prevent hypoglycemia. Because to their weakened immune systems, diabetic patients should be given antibiotics before undergoing tooth extraction or surgery, and they should continue taking the antibiotics for three to five days after the procedure.

Around a third of diabetes individuals experience xerostomia, which may be caused by a decreased salivary flow and an increased salivary glucose level. Diabetic sialadenitis refers to a concomitant, bilateral, non-tender swelling of the parotid glands. Increased vulnerability to opportunistic infections like *Candida albicans* is a consequence of xerostomia (dry mouth). In addition to this, the mouth can show signs of erythematous candidiasis, which manifests as central papillary atrophy of the dorsal tongue papillae, mucormycosis, benign migratory glossitis, changed taste, and burning mouth syndrome. In poorly controlled diabetes, high rates of dental caries are caused by elevated levels of glucose in the saliva and crevicular fluid.

Management of Oral Health: Dental care is tailored to each form of diabetes, as described below:

Non-insulin Dependent Patients	Insulin Controlled Patients	If Extensive Surgery Needed
- All dental procedures can be performed without special precautions, unless complication of diabetes present.	- All dental procedures can be performed. - Early morning appointments patients advised taking usual insulin dosage and normal meals on day of appointment. - If symptoms of insulin reaction occur, informed to dentist before procedure. - Source of glucose available and given to the patients if symptoms of insulin reaction occur.	- Dietary consultation required with physician during postoperative period. - Antibiotic prophylaxis can be considered for patients with brittle diabetes and those taking high dose of insulin which also have chronic status of oral infections.

- Crohn's Disease:

Lesions in either the intestine or the mouth do not appear to develop at the same rate or at the same time. Patients with this condition typically exhibit angular cheilitis, hyperplastic stiff mucosa, and cobble stoning of the buccal mucosa alongside generalised swelling of one or both lips. Other symptoms include gingival inflammation and cervical lymphadenopathy in addition to painful linear ulcerations in the buccal vestibule and painless localized swellings anywhere on the lips or face.

- Anemia:

Atrophic glossitis, mucosal pallor, and angular cheilitis are oral symptoms of iron deficiency anemia, the most prevalent hematologic condition. Dysphagia from pharyngoesophageal ulcerations is an oral symptom of Plummer Vinson Syndrome. Mandibular salmonella osteomyelitis causes osteoporosis and erosion, then osteosclerosis, and is one of the oral consequences encountered with sickle cell anemia.

Treatment options for anemia vary depending on the condition's severity and root cause:

1. Ferrous sulphate, ferrous fumarate, and ferrous gluconate are all iron supplements that can be used orally. In order to maximize the body's absorption of ingested iron, it's best to take these pills with a glass of orange juice or another source of vitamin C.
2. It is possible to boost erythropoiesis rates by administering injectable iron at the same time as erythropoietin.
3. Third, patients with hemoglobin levels below 60 to 80 g/L (6 to 8 g/dL) require transfusions.

4. Four, hyperbaric oxygen (HBO) is advised to provide oxygen into tissues when a patient's condition is critical and blood transfusion is not possible for medical or religious reasons. When there is a risk of infection or a blood type that cannot be transfused.
  5. The fifth type of drug is called erythropoiesis-stimulating agents (ESA), and it encourages the bone marrow to produce new RBCs (red blood cells). Some medical situations call for them, including severe anemia (kidney failure), major surgery, chemotherapy, and even some HIV/AIDS medications.
- Ulcerative Colitis:

It has been linked to destructive mouth ulcerations caused by immune-mediated vasculitis. These sores resemble aphthous ulcers, however they develop far less commonly than Crohn's lesions. Many painless intraepithelial micro abscesses connected in linear or serpentine tracks are characteristic of pyostomatitis vegetans (inflammatory stomatitis), which typically affects the labial mucosa, soft palate, and ventral tongue.

Administration in Dentistry: Preventative and routine dental care on a regular basis. The patient's capacity to undertake significant dental treatments is determined by an evaluation of hypothalamus, pituitary, and adrenal cortical function. Non-steroidal anti-inflammatory medicines (NSAIDs) should be avoided because they may cause an exacerbation; instead, paracetamol should be used. Topical steroid therapy has been shown to be effective in treating oral inflammatory and granulomatous lesions associated with inflammatory bowel diseases.

- Leukemia:

Gingival hypertrophy, petechiae, ecchymosis, mucosal ulcers, and bleeding are common oral consequences of leukaemia. Numb chin syndrome, also known as mental nerve neuropathy, is sometimes the first symptom experienced by those who have this condition. Mucormycosis of the nasal cavity and paranasal sinuses may be indicated by the presence of ulcerations and necrosis of the palatal mucosa. Septicemia can result from a bacterial infection of the mouth. Herpes simplex virus (HSV) reactivation leading to oral mucositis has been reported in patients undergoing chemotherapy for leukaemia. Oral mucositis is brought on by the chemotherapy medications because they thin the protective mucosal layer and inhibit the bone marrow, allowing the opportunistic organisms to infiltrate.

Dental Management: hematologic malignancies, as well as preventative measures and considerations in dentistry management, are as follows:

Prior to dental treatment	During dental treatment
1. Advised consultation with the specialist	1. Bleeding tendency
2. Blood investigation reports, detailed history, proper clinical and radiographic examination	2. Increased risk of infection.
3. Dental treatment should be performed before starting the chemotherapy or radiotherapy.	3. Risk of developing osteonecrosis of the jaw.
4. Long-term remission or maintenance phase patient can undergo dental treatment, while patients with advanced or relapsed disease with poor prognosis advised to receive palliative or urgent treatment only.	4. Anemia
	5. Corticosteroids treatment.
	6. Secondary malignancies.
	7. Specific Considerations.

- Pulmonary disease:

COPD Both chronic bronchitis and emphysema fall under the umbrella term of chronic obstructive pulmonary disease (COPD), which describes a group of pulmonary disorders defined by chronic, irreversible airflow limitation from the lungs. Most cases of chronic obstructive pulmonary disease (COPD) are brought on by years of exposure to lung irritants that cause permanent damage to the lungs and airways. The most important factor in developing COPD is smoking cigarettes. Recurrent respiratory tract infection, air pollution, cotton textile dust, inheritance, and age are also contributors to chronic obstructive pulmonary disease (COPD). A few of COPD's telltale symptoms are It's common parlance to refer to persistent coughs or those that generate lots of mucus as "smoker's cough." Experiencing difficulty breathing, particularly with exercise, Wheezing, Severe discomfort in the chest.

Unfortunately, COPD is not always the cause of such symptoms. Patients with severe COPD are typically prescribed bronchodilators like theophylline, inhaled beta agonists, or inhaled anticholinergics; in severe situations, patients are also given long acting medicines and inhaled corticosteroids or brief courses of systemic corticosteroids. Patients with COPD who are already on steroid treatment should have the option of supplemental care at the dental office before undergoing any significant procedures. It's best to stay away from drugs like sedatives, hypnotics, and narcotics that

slow your breathing. Dentist chairs don't always recline, so it's possible that keeping patients in an upright position can help them manage their excessive sputum and mucus. During operations, individuals with severe COPD should not receive more oxygen than normal unless their doctor specifically orders it.

- Hemophilia:

Problems with blood clotting might be caused by a lack of a certain coagulation factor, a problem with the platelets, a problem with the blood vessels, or a problem with the fibrinolytic system. Common congenital coagulation defect disorders include hemophilia A, which is caused by a lack of clotting factor VIII, hemophilia B, also called Christmas disease, caused by a lack of clotting factor IX, hemophilia C, also known as Rosenthal syndrome, caused by a lack of clotting factor XI, and von Willebrand's disease. 7 Very slow clotting times and a propensity for profuse bleeding are also potential dangers for persons with hemophilia.

Patient's factor VIII levels are increased, factor VIII is replaced, and fibrinolysis is inhibited as part of the management of hemophilia A prior to dental surgery.

Condition	Treatment and dose	Potential complications
Mild bleeding	Dose: 15 U/kg factor VIII every 8–12 hours for 1–2 days Target: 30% of normal level	Hemarthrosis, oropharyngeal or dental bleeding, epistaxis, hematuria
Major bleeding	Dose: 50 U/kg factor VIII every 8–12 hours for 7–14 days Target: 80% to 100%	Same potential complications as for mild bleeding, as well as CNS haemorrhage, retroperitoneal haemorrhage, gastrointestinal bleeding of normal level

Treatment for people with bleeding problems requires consideration of both the severity of the disorder and the extent of the dental operation being performed. Patients with moderate bleeding disorders may only need minor or no adjustments to the process, but in those with severe bleeding disorders, the primary goal is to restore the patient's hemostatic system to acceptable levels and maintain hemostasis via local and adjuvant approaches. Before beginning any kind of intrusive treatment, a doctor's opinion should be sought. Replacement of missing components is necessary in cases of irreversible coagulopathies. Nerve block injections are not recommended for hemophilic patients because the anaesthetic solution may cause a hematoma to form if deposited in a highly vascularized location. Gross swelling, pain, dysphasia, respiratory obstruction, and the danger of mortality from asphyxiation can result from blood extravasation in the oropharyngeal area due to either an inferior alveolar block or in the pterygoid plexus. Alternatives to nerve block with a vasoconstrictor (where possible) in such patients include anaesthetic infiltration and intraligamentary anaesthesia. When possible, hemophilia patients should undergo nonsurgical endodontic therapy rather than extraction.

- Thyroid patients:

Thyroid hormones, including thyroxine (T4) and triiodothyronine, are secreted by the thyroid gland, an endocrine gland in the neck (T3). The body's metabolic rate is controlled by thyroid hormones, which also aid in proper brain growth and function. Fat, protein, and carbohydrate metabolism are all controlled by these hormones. Thyroid hormones have a crucial role in controlling a wide variety of physiological processes, including body temperature, metabolism, and energy expenditure, among others. The degree of abnormal thyroid function determines whether a patient is considered euthyroid, hypothyroid, or hyperthyroid. Hyperthyroidism is characterized by abdominal pain, hives, and rapid heart rate, while hypothyroidism is characterized by anemia, cardiomegaly, cold intolerance, constipation, cretinism (in children), dry hair, elevated creatine, goiter, hyperlipidemia, hypertelorism, hypotension, inverted T waves in electrocardiogram, lethargy, low amplitude QRS waves in electrocardiogram, myxedema, paresthesia Avoid surgical procedures, treat oral infection, and avoid CNS depressants like narcotics and barbiturates; in well-controlled cases, avoid oral infections and implement normal procedures and management; in the extremely rare event of a medical emergency, recognize the symptoms of myxedematous coma, seek medical assistance,

administer 100 milligrams of hydrocortisone, and perform cardiopulmonary resuscitation. Avoiding adrenaline and preventing the transmission of infection are two key components of dental treatment for hyperthyroidism. If a thyrotoxic crisis develops, treatment must be stopped. Because emotional or physical stress, or even surgical stress, can precipitate a thyrotoxic crisis, it is crucial that these patients have their stress levels managed. Propylthiouracil, like other anti-thyroid medicines, has anti-vitamin K activity and can cause hypoprothrombinemia and bleeding, which offers a risk for hemorrhage; as a result, patients using PTU need to be carefully reviewed before undergoing surgery or invasive dental treatment.

- Cardiac problems:

In addition to increasing the likelihood that you will experience cardiovascular disease like angina, myocardial infarction, and cerebrovascular events like stroke, the high incidence and prevalence of hypertension in the general population makes it a significant health issue. National Committee on the Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (JNC 7) coined the term "pre-hypertension" to describe people with systolic blood pressure of 120-139 mmHg or with diastolic blood pressure of 80-89 mmHg. The consensus values for normal blood pressure are under 90 mmHg for diastolic pressure and under 140 mmHg for systolic pressure. Patients with hypertension should be instructed to continue taking their prescribed medications on the day of dental treatment, since this condition does not pose a risk in clinical practise when under tight control. The patient's blood pressure should be taken before any dental work is done, and if it's too high, the procedure will have to be delayed. Morning appointments at the dentist are preferable because they tend to be shorter. Patients with anxiety should take an anxiolytic medication, such as diazepam (5-10 mg), the night before and again within one to two hours before their dental consultation. Use of two anaesthetic cartridges (1.8mlx2) with vasoconstrictor is the absolute maximum that should be used for a local anaesthetic, and intravascular injection should be avoided. A greater dose of anaesthetic may be administered if necessary, but without the use of a vasoconstrictor. Adrenaline should not be combined with absorbable sutures. If the patient's blood pressure is not under control, he should be referred to a doctor. For dental emergencies, it's best to take a conservative approach that focuses on relieving pain and preventing infection. Once blood pressure is under control, surgical intervention should be postponed. Ibuprofen, indomethacin, and naproxen are nonsteroidal anti-inflammatory medicines (NSAIDs) that may interact with antihypertensive therapies (beta-blockers, diuretics, ACEIs) and reduce the efficacy of these medications in treating hypertension. Drug interactions typically involve administration of both medications for more than five days; hence, NSAIDs should not be recommended for longer than this five-day period. It is recommended to provide furosemide to patients with severe hypertension (>120/210 mmHg) (40 mg, via the oral route). When this doesn't lower blood pressure enough, captopril is the next step (25 mg via the oral or sublingual route). Refer the patient to the nearest hospital if the blood pressure does not start to drop within 30 minutes after taking these measures.

Ischemic heart disease (HD) is characterized by a decrease in blood flow to the heart muscle as a result of plaque buildup in the coronary arteries. Myocardial infarction, unstable angina, and sudden cardiac death are all examples of ischemic heart disease.

Patients receiving anti-platelet or anti-coagulant treatment for a myocardial infarction (MI) or angina should have their CT, BT, PT, and pTT/INR monitored prior to any surgical operation or dental extractions. Moreover, the appointments ought to be shorter and stress management techniques ought to be implemented.

- Epilepsy:

The dental office can trigger seizures in epileptic patients, even those with good seizure control. Carbamazepine, phenobarbitone, sodium valproate, phenytoin, lamotrigine, vigabatrin, and gabapentin are all anticonvulsants that can be used as preventative measures. Stop the dental process and have the patient lie down in a supine position; provide adequate protection for the patient during an attack; In the case of LA-induced convulsions, continued maintenance of a patent airway and adequate oxygenation should be prioritized until the LA level in the brain drops below the seizure threshold. The anticonvulsant diazepam is given intravenously (IV) at a dose of 0.2-0.5mg/kg every 3-5 minutes.

- Renal problems:

There must be close communication between the dentist and the treating nephrologist when a patient has undergone a kidney transplant and needs dental therapy. Following a transplant, patients are typically treated with a cocktail of tacrolimus, mycophenolate, and prednisolone and other immunosuppressants to reduce the likelihood of organ rejection. Other medications like as cyclosporine, sirolimus, and azathioprine are used instead in rare cases. Due to the risk of bleeding that persists even after a kidney transplant, as well as the possibility of coumarin anticoagulants being used in the minority of patients who also suffer from cardiovascular disease, a recent INR should be required

before any dental work that may involve bleeding. If the INR is greater than 2.5, the patient should consult with a nephrologist.

To improve the oral and overall health of patients with renal failure (on dialysis), dental and medical specialists must work closely together in the management of the patient's dental care. A full blood count and coagulation testing are required before undergoing any invasive dental treatment. Antibiotic prophylaxis should be considered for any oral infection if bleeding and/or septicemia are imminent concerns. Constantly checking your blood pressure is a good idea. Antibiotics such as penicillins, clindamycin, and cephalosporins can be given at the regular dosages and are recommended, however the dosing interval should be lengthened. The painkiller of preference is paracetamol (PCM), while aspirin should be avoided because to its antiplatelet action. Heparin has a half-life of around four hours, therefore dental work should be scheduled on days when the patient is not receiving dialysis. In order to prevent fatal bleeding in people with renal insufficiency, desmopressin has been advocated. Rinses containing tranexamic acid or oral administration of 10-15 mg/kg body weight daily, split into 2-3 doses, may also be effective.

- Infectious diseases:

HIV is a retrovirus that causes AIDS, which is an immune deficiency disease (human immunodeficiency virus). Sexual contact between an infected person and someone who is not affected, sharing needles or other injecting equipment, and becoming pregnant, giving birth, or breastfeeding are all ways that HIV can spread to a new host. The blood of an infected patient can potentially cause infection if it comes into contact with the skin or mucous membranes.

AIDS treatment typically focuses on three main pillars: (1) antiretroviral therapy (ART), (2) opportunistic infection prophylaxis, and (3) management of HIV-related complications. All new patients should undergo a thorough examination that includes a medical history review, a head and neck examination, an intraoral soft tissue examination, and a dental and periodontal examination. Nutritional support is essential for cancer patients and other immunocompromised individuals. As immunosuppression is a major role in the development of oral lesions in AIDS patients, it is crucial that these patients be treated with analgesics or drugs specifically designed to address oral lesions. Many doctors don't bother to do in-depth intraoral examinations on normal patients, therefore getting a good oral exam is crucial. Dentists have a duty to check for oral cancer and any intraoral lesions that could be indicators of HIV, oral illness, or the oral/perioral manifestations of other systemic disorders. Kaposi's sarcoma, facial palsy, trigeminal neuropathy, recurrent thrombocytopenic purpura, recurrent aphthous ulcers, salivary gland enlargement, xerostomia that may lead to dental caries, methicillin-resistant strep throat, and thrush are all oral manifestations of AIDS. Dentists have a responsibility to do thorough oral cancer screenings and address any issues that are found. Moreover, before performing any invasive operation, it is best to speak with the patient's primary care physician to assess the patient's current viral load and CD4 count. A dentist who is concerned about his health should adopt chemo-prophylactic measures in case of exposure, in addition to practising universal precautions.

#### **Conclusion:**

It is clear from the foregoing that it is crucial for dental OPD staff to be well-versed in the medical and dental treatment of a wide range of medical illnesses. The patient's life may be in jeopardy if even a brief period of stress triggers an angina attack, or if the medical history is not taken with the utmost care and precision. To effectively manage such patients and to forestall medical problems, dentists must conduct thorough medical histories and physical examinations.

Diseases that affect the entire body frequently show their effects orally first. A familiarity with systemic disorders is crucial for effective clinical work. Given its accessibility for visual study and assessment by palpation, the mouth provides a portal for easy observation of signs and symptoms of many systemic disorders. Primary care doctors need to be familiar with the oral signs of systemic disease and be able to spot the accompanying changes in their patients in order to provide the best possible care.



### References:

- Diz, P., Scully, C., & Sanz, M. (2013). Dental implants in the medically compromised patient. *Journal of dentistry*, 41(3), 195-206.
- Dougall, A., Hayes, M., & Daly, B. (2017). A systematic review of the use of local analgesia in medically compromised children and adolescents. *European Archives of Paediatric Dentistry*, 18, 331-343.
- Kühnisch, J., Daubländer, M., Klingberg, G., Dougall, A., Spyridonos Loizides, M., Stratigaki, E., ... & Gizani, S. (2017). Best clinical practice guidance for local analgesia in paediatric dentistry: an EAPD policy document. *European Archives of Paediatric Dentistry*, 18, 313-321.
- Alqahtani, H. (2019). Medically compromised patients in orthodontic practice: review of evidence and recommendations. *International Orthodontics*, 17(4), 776-788.
- Lam, O. L., Bandara, H. M. H. N., Samaranayake, L. P., McGrath, C., & Li, L. S. (2012). Oral health promotion interventions on oral yeast in hospitalised and medically compromised patients: a systematic review. *Mycoses*, 55(2), 123-142.
- Scully, C., Hobkirk, J., & D DIOS, P. (2007). Dental endosseous implants in the medically compromised patient 1. *Journal of oral rehabilitation*, 34(8), 590-599.
- Fleming, P. (2012). Medically Compromised Patients. *Clinical Cases in Pediatric Dentistry*, 1, 3.
- Pimentel, E. L. C., Azevedo, V. M. P., Castro, R. D. A. L., Reis, L. C., & Lorenzo, A. D. (2013). Caries experience in young children with congenital heart disease in a developing country. *Brazilian oral research*, 27, 103-108.
- 
- Yadav P, Gupta ND, Tewari RK, Agarwal N, Sinha A, Arora D. (2016). Oral health status of asthmatic children: a review. *European Journal of Pharmaceutical and Medical Research*; 3(3): 192-197.
- Inamdar P, Verma P. Endocrine and Metabolic Disorders. In: Gupta PV, editor. (2012). *Understanding and Management of Special child in Pediatric Dentistry*, 1st ed. New Delhi: Jaypee Brothers Medical Publisher. p. 212-220
- Van Meter, K. W. (2005). A systematic review of the application of hyperbaric oxygen in the treatment of severe anemia: an evidence-based approach. *Undersea Hyperb Med*, 32(1), 61-83.
- Fabuel, L. C., Soriano, Y. J., & Pérez, M. G. S. (2010). Dental management of patients with endocrine disorders. *Journal of Clinical and Experimental Dentistry*, 2(4), 196-203.
- Rahman, S. S., Faruque, M., Khan, M. H. A., & Hossain, S. A. (2011). Dental management of COPD patient. *Bangladesh Medical Journal Khulna*, 44(1-2), 21-24.
- Nazif, M. (1970). Local anesthesia for patients with hemophilia. *ASDC Journal of Dentistry for Children*, 37(1), 79-84.
- Leatherdale, R. A. (1960). Respiratory obstruction in haemophilic patients. *British Medical Journal*, 1(5182), 1316.
- Little, J. W. (2006). Thyroid disorders. Part II: hypothyroidism and thyroiditis. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, and Endodontology*, 102(2), 148-153.
- Joint National Committee on Detection, Evaluation, Treatment of High Blood Pressure, & National High Blood Pressure Education Program. Coordinating Committee. (1995). *Report of the Joint National Committee on Detection, Evaluation, and Treatment of High Blood Pressure*. National Heart, Lung, and Blood Institute, National High Blood Pressure Education Program.
- Fatima, D., Wahengbam, P., Verma, S., Agnihotri, A., & Sehrawat, M. (2021). A brief review in dental management of medically compromised patients. *IP Annals of Prosthodontics and Restorative Dentistry*, 7(01), 5-11.